



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

THE MEGALOPYGID GENUS *TROSIA*, WITH DESCRIPTION OF A NEW SPECIES.

BY HARRISON G. DYAR.

Genus *Trosia* Hübner.

1822—*Trosia* HÜBNER, Verz. bek. Schmett. 196.

1855—*Sciathos* WALKER, Cat. Brit. Mus. III, 752.

1856—*Edebea* WALKER, Cat. Brit. Mus. VII, 1755.

1874—*Isochroma* FELDER, Reise Novara, Lep. IV, pl. 83.

Kirby calls this genus *Sciathos* in the catalogue, omitting Hübner's term. Druce and others also neglect *Trosia*, perhaps from a prejudice against the Verzeichniss names. In the following, species succeeded by a dash are unknown to me, except by description and are referred to this genus on the authority of the authors quoted.

SYNOPSIS OF SPECIES.

Thorax and abdomen discolorous, thorax in part white, abdomen bright red.

Thorax without black marks.

Costa broadly red ; wings in part pure white, at least in a stripe next the red margin.....1. **tricolora** Fab.

Costa very narrowly red ; wings without any pure white.

2. **obsolescens** Dyar

Thorax with a black band in front and on each side. 3. **purens** Walk.

Thorax and abdomen concolorous, abdomen not bright red.

Costal margin red ; a transverse row of black dots. 4. **diamas** Cram.

Fore wings without marks, uniformly ochraceous.....5. **ribbei** Druce.

1. *T. tricolora* Fab.

1787—*Bombyx tricolora* FABRICIUS, Mant. Ins. II, 114.

1790—*Bombyx punctigera* STOLL, Pap. Exot. suppl. pl. 34, fig. 1 (nec Linn.).

1822—*Trosia tricolora* HÜBNER, Verz. bek. Schmett. 196.

1855—*Sciathos punctiger* WALKER, Cat. Brit. Mus. 752.

1856—*Sciathos punctiger* WALKER, Cat. Brit. Mus. VII, 1711.

1874—*Isochroma fallax* FELDER, Reise Novara, pl. 83, figs. 18, 19.

1887—*Sciathos punctigera* DRUCE, Biol. Cent.-Am. Lep. I, 212.

1892—*Sciathos punctigera* KIRBY, Cat. Lep. Het. I, 540.

Kirby makes this the *punctigera* of Linnæus (Syst. Nat., 509, no. 67), but the description does not coincide in the least with this species, either in structure, color or habitat.

2. *T. obsolescens*, sp. nov.

Head white in front, red on the vertex behind the antennæ and shading to red below

on the palpi. Thorax white above with six clusters of red hairs; abdomen red, white at tip. Fore wings uniformly pale ochreous, almost white, appearing pinkish from the red scales below; costa very narrowly red at base, dark ochre at apical portion. A straight row of eight small black spots between the veins beyond the middle of the wing, the sixth spot between veins 4 and 5, the seventh opposite the discal cross-vein and the eighth between vein 6 and the stalk of veins 7 to 10. Hind wings red-fringe ochreous. Below both wings as secondaries above; body largely white; coxæ and femora red above, tibiae and tarsi ringed with black. Expanse, 27 mm.

Nearly allied to *T. tricolora* Fab., which is however an inhabitant of the tropical regions, whereas this comes from the Mexican plateau.

One male, Nogales, Koebele collector, August 15, 1898, U. S. Nat. Mus., type no. 4104. Nogales is a town on the border line between Arizona and Mexico.

3. *T. purens* Walk.

1856—*Edebea purens* WALKER, Cat. Brit. Mus. VII, 1755.

1892—*Sciathos purens* KIRBY, Cat. Lep. Het. I, 540.

Sir G. F. Hampson has kindly examined Walker's types of the species for me and the generic characters correspond with *Trosia*.

4. *T. dimas* Cram. —

1775—*Bombyx dimas* CRAMER, Pap. Exot. I, pl. 59 C.

1822—*Trosia dimas* HÜBNER, Verz. bek. Schmett. 196.

1854—*Chrysauge dimas* WALKER, Cat. Brit. Mus. II, 375.

1892—*Idalus* (?) *dimas* KIRBY, Cat. Lep. Het. I, 198.

1894—*Sciathos dimas* DOGNIN, Lep. Loja. 173.

1897—*Sciathos dimas* DRUCE, Biol. Cent.-Am., Lep. Het., II, 440.

5. *T. ribbei* Druce. —

1898—*Sciathos ribbei* DRUCE, Biol. Cent.-Am. Lep. Het. II, 441, pl. 88, fig. 1.

NEW SPECIES OF SYNTOMIDÆ.

BY HARRISON G. DYAR.

Pseudapinconoma elegans Auriv. var. *curriei*, var. nov.

Under side of thorax entirely crimson, legs white, femora and basal half of hind tibiae crimson above; abdomen bluish gray, segmental black bands linear, the basal segments with orange hair and the lateral tufts orange; a dorsal series of crimson dots. Wings as in *elegans*, but the hyaline patches between veins 2 and 6 large and diffuse, reaching nearly to the termen, with ill defined outer border.

Two males, Mt. Coffee, Liberia (R. P. Currie). U. S. Nat. Mus., type no. 4247.